

IMPORTANT SAFETY INFORMATION

Serious Adverse Reactions

Please refer to the full Prescribing Information for important dose management information specific to adverse reactions.

- **Immune-mediated pneumonitis.** Immune-mediated pneumonitis or interstitial lung disease, including fatal cases, have occurred. Permanently discontinue TECENTRIQ for Grade 3 or 4 pneumonitis
- **Immune-mediated hepatitis.** Immune-mediated hepatitis and liver test abnormalities, including fatal cases, have occurred. Permanently discontinue TECENTRIQ for AST or ALT elevations more than 8 times the upper limit of normal or total bilirubin more than 3 times the upper limit of normal
- **Immune-mediated colitis.** Immune-mediated diarrhea or colitis have occurred. Permanently discontinue TECENTRIQ for Grade 4 diarrhea or colitis
- **Immune-mediated endocrinopathies.** Thyroid disorders, adrenal insufficiency, type 1 diabetes mellitus, including diabetic ketoacidosis, and hypophysitis/hypopituitarism have occurred
- **Other immune-mediated adverse reactions.** TECENTRIQ can cause severe and fatal immune-mediated adverse reactions. These immune-mediated reactions may involve any organ system. Permanently discontinue TECENTRIQ for Grade 4 immune-mediated adverse reactions involving a major organ
- **Infections.** Severe infections, including fatal cases, have occurred
- **Infusion-related reactions.** Severe or life-threatening infusion-related reactions have occurred. Permanently discontinue TECENTRIQ in patients with Grade 3 or 4 infusion-related reactions
- **Embryo-fetal toxicity.** TECENTRIQ can cause fetal harm in pregnant women. Verify pregnancy status prior to initiating TECENTRIQ. Advise females of reproductive potential of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with TECENTRIQ and for at least 5 months after the last dose
- Advise female patients not to breastfeed while taking TECENTRIQ and for at least 5 months after the last dose

Most Common Adverse Reactions

The most common adverse reactions (rate $\geq 20\%$) in patients who received TECENTRIQ in combination with bevacizumab for HCC were hypertension (30%), fatigue/asthenia (26%), and proteinuria (20%).

You may report side effects to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. You may also report side effects to Genentech at 1-888-835-2555.

Please see accompanying full Prescribing Information for additional Important Safety Information.